



# Highlights of Changes from DSM-IV-TR to DSM-5



Changes made to the DSM-5 diagnostic criteria and texts are outlined in this chapter in the same order in which they appear in the DSM-5 classification. This is not an exhaustive guide; minor changes in text or wording made for clarity are not described here. It should also be noted that Section I of DSM-5 contains a description of changes pertaining to the chapter organization in DSM-5, the multiaxial system, and the introduction of dimensional assessments (in Section III).

## Terminology

The phrase “general medical condition” is replaced in DSM-5 with “another medical condition” where relevant across all disorders.

## Neurodevelopmental Disorders

### Intellectual Disability (Intellectual Developmental Disorder)

Diagnostic criteria for intellectual disability (intellectual developmental disorder) emphasize the need for an assessment of both cognitive capacity (IQ) and adaptive functioning. Severity is determined by adaptive functioning rather than IQ score. The term mental retardation was used in DSM-IV. However, *intellectual disability* is the term that has come into common use over the past two decades among medical, educational, and other professionals, and by the lay public and advocacy groups. Moreover, a federal statute in the United States (Public Law 111-256, Rosa’s Law) replaces the term “mental retardation with intellectual disability. Despite the name change, the deficits in cognitive capacity beginning in the developmental period, with the accompanying diagnostic criteria, are considered to constitute a mental disorder. The term *intellectual developmental disorder* was placed in parentheses to reflect the World Health Organization’s classification system, which lists “disorders” in the International Classification of Diseases (ICD; ICD-11 to be released in 2015) and bases all “disabilities” on the International Classification of Functioning, Disability, and Health (ICF). Because the ICD-11 will not be adopted for several years, *intellectual disability* was chosen as the current preferred term with the bridge term for the future in parentheses.

### Communication Disorders

The DSM-5 communication disorders include language disorder (which combines DSM-IV expressive and mixed receptive-expressive language disorders), speech sound disorder (a new name for phonological disorder), and childhood-onset fluency disorder (a new name for stuttering). Also included is social (pragmatic) communication disorder, a new condition for persistent difficulties in the social uses of verbal and nonverbal communication. Because social communication deficits are one component of autism spectrum disorder (ASD), it is important to note that social (pragmatic) communication disorder cannot be diagnosed in the presence of restricted repetitive behaviors, interests, and activities (the other component of ASD). The symptoms of some patients diagnosed with DSM-IV pervasive developmental disorder not otherwise specified may meet the DSM-5 criteria for social communication disorder.

### Autism Spectrum Disorder

Autism spectrum disorder is a new DSM-5 name that reflects a scientific consensus that four previously separate disorders are actually a single condition with different levels of symptom severity in two core

domains. ASD now encompasses the previous DSM-IV autistic disorder (autism), Asperger's disorder, childhood disintegrative disorder, and pervasive developmental disorder not otherwise specified. ASD is characterized by 1) deficits in social communication and social interaction and 2) restricted repetitive behaviors, interests, and activities (RRBs). Because both components are required for diagnosis of ASD, social communication disorder is diagnosed if no RRBs are present.

### **Attention-Deficit/Hyperactivity Disorder**

The diagnostic criteria for attention-deficit/hyperactivity disorder (ADHD) in DSM-5 are similar to those in DSM-IV. The same 18 symptoms are used as in DSM-IV, and continue to be divided into two symptom domains (inattention and hyperactivity/impulsivity), of which at least six symptoms in one domain are required for diagnosis. However, several changes have been made in DSM-5: 1) examples have been added to the criterion items to facilitate application across the life span; 2) the cross-situational requirement has been strengthened to "several" symptoms in each setting; 3) the onset criterion has been changed from "symptoms that caused impairment were present before age 7 years" to "several inattentive or hyperactive-impulsive symptoms were present prior to age 12"; 4) subtypes have been replaced with presentation specifiers that map directly to the prior subtypes; 5) a comorbid diagnosis with autism spectrum disorder is now allowed; and 6) a symptom threshold change has been made for adults, to reflect their substantial evidence of clinically significant ADHD impairment, with the cutoff for ADHD of five symptoms, instead of six required for younger persons, both for inattention and for hyperactivity and impulsivity. Finally, ADHD was placed in the neurodevelopmental disorders chapter to reflect brain developmental correlates with ADHD and the DSM-5 decision to eliminate the DSM-IV chapter that includes all diagnoses usually first made in infancy, childhood, or adolescence.

### **Specific Learning Disorder**

Specific learning disorder combines the DSM-IV diagnoses of reading disorder, mathematics disorder, disorder of written expression, and learning disorder not otherwise specified. Because learning deficits in the areas of reading, written expression, and mathematics commonly occur together, coded specifiers for the deficit types in each area are included. The text acknowledges that specific types of reading deficits are described internationally in various ways as dyslexia and specific types of mathematics deficits as dyscalculia.

### **Motor Disorders**

The following motor disorders are included in the DSM-5 neurodevelopmental disorders chapter: developmental coordination disorder, stereotypic movement disorder, Tourette's disorder, persistent (chronic) motor or vocal tic disorder, provisional tic disorder, other specified tic disorder, and unspecified tic disorder. The tic criteria have been standardized across all of these disorders in this chapter. Stereotypic movement disorder has been more clearly differentiated from body-focused repetitive behavior disorders that are in the DSM-5 obsessive-compulsive disorder chapter.

## **Schizophrenia Spectrum and Other Psychotic Disorders**

### **Schizophrenia**

Two changes were made to DSM-IV Criterion A for schizophrenia. The first change is the elimination of the special attribution of bizarre delusions and Schneiderian first-rank auditory hallucinations (e.g., two or more voices conversing). In DSM-IV, only one such symptom was needed to meet the diagnostic requirement for Criterion A, instead of two of the other listed symptoms. This special attribution was

removed due to the nonspecificity of Schneiderian symptoms and the poor reliability in distinguishing bizarre from nonbizarre delusions. Therefore, in DSM-5, two Criterion A symptoms are required for any diagnosis of schizophrenia. The second change is the addition of a requirement in Criterion A that the individual must have at least one of these three symptoms: delusions, hallucinations, and disorganized speech. At least one of these core “positive symptoms” is necessary for a reliable diagnosis of schizophrenia.

### **Schizophrenia subtypes**

The DSM-IV subtypes of schizophrenia (i.e., paranoid, disorganized, catatonic, undifferentiated, and residual types) are eliminated due to their limited diagnostic stability, low reliability, and poor validity. These subtypes also have not been shown to exhibit distinctive patterns of treatment response or longitudinal course. Instead, a dimensional approach to rating severity for the core symptoms of schizophrenia is included in Section III to capture the important heterogeneity in symptom type and severity expressed across individuals with psychotic disorders.

### **Schizoaffective Disorder**

The primary change to schizoaffective disorder is the requirement that a major mood episode be present for a majority of the disorder’s total duration after Criterion A has been met. This change was made on both conceptual and psychometric grounds. It makes schizoaffective disorder a longitudinal instead of a cross-sectional diagnosis—more comparable to schizophrenia, bipolar disorder, and major depressive disorder, which are bridged by this condition. The change was also made to improve the reliability, diagnostic stability, and validity of this disorder, while recognizing that the characterization of patients with both psychotic and mood symptoms, either concurrently or at different points in their illness, has been a clinical challenge.

### **Delusional Disorder**

Criterion A for delusional disorder no longer has the requirement that the delusions must be non-bizarre. A specifier for bizarre type delusions provides continuity with DSM-IV. The demarcation of delusional disorder from psychotic variants of obsessive-compulsive disorder and body dysmorphic disorder is explicitly noted with a new exclusion criterion, which states that the symptoms must not be better explained by conditions such as obsessive-compulsive or body dysmorphic disorder with absent insight/delusional beliefs. DSM-5 no longer separates delusional disorder from shared delusional disorder. If criteria are met for delusional disorder then that diagnosis is made. If the diagnosis cannot be made but shared beliefs are present, then the diagnosis “other specified schizophrenia spectrum and other psychotic disorder” is used.

### **Catatonia**

The same criteria are used to diagnose catatonia whether the context is a psychotic, bipolar, depressive, or other medical disorder, or an unidentified medical condition. In DSM-IV, two out of five symptom clusters were required if the context was a psychotic or mood disorder, whereas only one symptom cluster was needed if the context was a general medical condition. In DSM-5, all contexts require three catatonic symptoms (from a total of 12 characteristic symptoms). In DSM-5, catatonia may be diagnosed as a specifier for depressive, bipolar, and psychotic disorders; as a separate diagnosis in the context of another medical condition; or as an other specified diagnosis.

## Bipolar and Related Disorders

### Bipolar Disorders

To enhance the accuracy of diagnosis and facilitate earlier detection in clinical settings, Criterion A for manic and hypomanic episodes now includes an emphasis on changes in activity and energy as well as mood. The DSM-IV diagnosis of bipolar I disorder, mixed episode, requiring that the individual simultaneously meet full criteria for both mania and major depressive episode, has been removed. Instead, a new specifier, “with mixed features,” has been added that can be applied to episodes of mania or hypomania when depressive features are present, and to episodes of depression in the context of major depressive disorder or bipolar disorder when features of mania/hypomania are present.

### Other Specified Bipolar and Related Disorder

DSM-5 allows the specification of particular conditions for other specified bipolar and related disorder, including categorization for individuals with a past history of a major depressive disorder who meet all criteria for hypomania except the duration criterion (i.e., at least 4 consecutive days). A second condition constituting an other specified bipolar and related disorder is that too few symptoms of hypomania are present to meet criteria for the full bipolar II syndrome, although the duration is sufficient at 4 or more days.

### Anxious Distress Specifier

In the chapter on bipolar and related disorders and the chapter on depressive disorders, a specifier for anxious distress is delineated. This specifier is intended to identify patients with anxiety symptoms that are not part of the bipolar diagnostic criteria.

## Depressive Disorders

DSM-5 contains several new depressive disorders, including disruptive mood dysregulation disorder and premenstrual dysphoric disorder. To address concerns about potential overdiagnosis and overtreatment of bipolar disorder in children, a new diagnosis, disruptive mood dysregulation disorder, is included for children up to age 18 years who exhibit persistent irritability and frequent episodes of extreme behavioral dyscontrol. Based on strong scientific evidence, premenstrual dysphoric disorder has been moved from DSM-IV Appendix B, “Criteria Sets and Axes Provided for Further Study,” to the main body of DSM-5. Finally, DSM-5 conceptualizes chronic forms of depression in a somewhat modified way. What was referred to as dysthymia in DSM-IV now falls under the category of persistent depressive disorder, which includes both chronic major depressive disorder and the previous dysthymic disorder. An inability to find scientifically meaningful differences between these two conditions led to their combination with specifiers included to identify different pathways to the diagnosis and to provide continuity with DSM-IV.

### Major Depressive Disorder

Neither the core criterion symptoms applied to the diagnosis of major depressive episode nor the requisite duration of at least 2 weeks has changed from DSM-IV. Criterion A for a major depressive episode in DSM-5 is identical to that of DSM-IV, as is the requirement for clinically significant distress or impairment in social, occupational, or other important areas of life, although this is now listed as Criterion B rather than Criterion C. The coexistence within a major depressive episode of at least three manic symptoms (insufficient to satisfy criteria for a manic episode) is now acknowledged by the specifier “with mixed features.” The presence of mixed features in an episode of major depressive disorder in-

creases the likelihood that the illness exists in a bipolar spectrum; however, if the individual concerned has never met criteria for a manic or hypomanic episode, the diagnosis of major depressive disorder is retained.

### **Bereavement Exclusion**

In DSM-IV, there was an exclusion criterion for a major depressive episode that was applied to depressive symptoms lasting less than 2 months following the death of a loved one (i.e., the bereavement exclusion). This exclusion is omitted in DSM-5 for several reasons. The first is to remove the implication that bereavement typically lasts only 2 months when both physicians and grief counselors recognize that the duration is more commonly 1–2 years. Second, bereavement is recognized as a severe psychosocial stressor that can precipitate a major depressive episode in a vulnerable individual, generally beginning soon after the loss. When major depressive disorder occurs in the context of bereavement, it adds an additional risk for suffering, feelings of worthlessness, suicidal ideation, poorer somatic health, worse interpersonal and work functioning, and an increased risk for persistent complex bereavement disorder, which is now described with explicit criteria in Conditions for Further Study in DSM-5 Section III. Third, bereavement-related major depression is most likely to occur in individuals with past personal and family histories of major depressive episodes. It is genetically influenced and is associated with similar personality characteristics, patterns of comorbidity, and risks of chronicity and/or recurrence as non-bereavement-related major depressive episodes. Finally, the depressive symptoms associated with bereavement-related depression respond to the same psychosocial and medication treatments as non-bereavement-related depression. In the criteria for major depressive disorder, a detailed footnote has replaced the more simplistic DSM-IV exclusion to aid clinicians in making the critical distinction between the symptoms characteristic of bereavement and those of a major depressive episode. Thus, although most people experiencing the loss of a loved one experience bereavement without developing a major depressive episode, evidence does not support the separation of loss of a loved one from other stressors in terms of its likelihood of precipitating a major depressive episode or the relative likelihood that the symptoms will remit spontaneously.

### **Specifiers for Depressive Disorders**

Suicidality represents a critical concern in psychiatry. Thus, the clinician is given guidance on assessment of suicidal thinking, plans, and the presence of other risk factors in order to make a determination of the prominence of suicide prevention in treatment planning for a given individual. A new specifier to indicate the presence of mixed symptoms has been added across both the bipolar and the depressive disorders, allowing for the possibility of manic features in individuals with a diagnosis of unipolar depression. A substantial body of research conducted over the last two decades points to the importance of anxiety as relevant to prognosis and treatment decision making. The “with anxious distress” specifier gives the clinician an opportunity to rate the severity of anxious distress in all individuals with bipolar or depressive disorders.

## **Anxiety Disorders**

The DSM-5 chapter on anxiety disorder no longer includes obsessive-compulsive disorder (which is included with the obsessive-compulsive and related disorders) or posttraumatic stress disorder and acute stress disorder (which is included with the trauma- and stressor-related disorders). However, the sequential order of these chapters in DSM-5 reflects the close relationships among them.

### **Agoraphobia, Specific Phobia, and Social Anxiety Disorder (Social Phobia)**

Changes in criteria for agoraphobia, specific phobia, and social anxiety disorder (social phobia) include deletion of the requirement that individuals over age 18 years recognize that their anxiety is excessive or unreasonable. This change is based on evidence that individuals with such disorders often overestimate the danger in “phobic” situations and that older individuals often misattribute “phobic” fears to aging. Instead, the anxiety must be out of proportion to the actual danger or threat in the situation, after taking cultural contextual factors into account. In addition, the 6-month duration, which was limited to individuals under age 18 in DSM-IV, is now extended to all ages. This change is intended to minimize overdiagnosis of transient fears.

### **Panic Attack**

The essential features of panic attacks remain unchanged, although the complicated DSM-IV terminology for describing different types of panic attacks (i.e., situationally bound/cued, situationally predisposed, and unexpected/uncued) is replaced with the terms unexpected and expected panic attacks. Panic attacks function as a marker and prognostic factor for severity of diagnosis, course, and comorbidity across an array of disorders, including but not limited to anxiety disorders. Hence, panic attack can be listed as a specifier that is applicable to all DSM-5 disorders.

### **Panic Disorder and Agoraphobia**

Panic disorder and agoraphobia are unlinked in DSM-5. Thus, the former DSM-IV diagnoses of panic disorder with agoraphobia, panic disorder without agoraphobia, and agoraphobia without history of panic disorder are now replaced by two diagnoses, panic disorder and agoraphobia, each with separate criteria. The co-occurrence of panic disorder and agoraphobia is now coded with two diagnoses. This change recognizes that a substantial number of individuals with agoraphobia do not experience panic symptoms. The diagnostic criteria for agoraphobia are derived from the DSM-IV descriptors for agoraphobia, although endorsement of fears from two or more agoraphobia situations is now required, because this is a robust means for distinguishing agoraphobia from specific phobias. Also, the criteria for agoraphobia are extended to be consistent with criteria sets for other anxiety disorders (e.g., clinician judgment of the fears as being out of proportion to the actual danger in the situation, with a typical duration of 6 months or more).

### **Specific Phobia**

The core features of specific phobia remain the same, but there is no longer a requirement that individuals over age 18 years must recognize that their fear and anxiety are excessive or unreasonable, and the duration requirement (“typically lasting for 6 months or more”) now applies to all ages. Although they are now referred to as specifiers, the different types of specific phobia have essentially remained unchanged.

### **Social Anxiety Disorder (Social Phobia)**

The essential features of social anxiety disorder (social phobia) (formerly called social phobia) remain the same. However, a number of changes have been made, including deletion of the requirement that individuals over age 18 years must recognize that their fear or anxiety is excessive or unreasonable, and duration criterion of “typically lasting for 6 months or more” is now required for all ages. A more significant change is that the “generalized” specifier has been deleted and replaced with a “performance only” specifier. The DSM-IV generalized specifier was problematic in that “fears include most social situations” was difficult to operationalize. Individuals who fear only performance situations (i.e., speaking

or performing in front of an audience) appear to represent a distinct subset of social anxiety disorder in terms of etiology, age at onset, physiological response, and treatment response.

### **Separation Anxiety Disorder**

Although in DSM-IV, separation anxiety disorder was classified in the section “Disorders Usually First Diagnosed in Infancy, Childhood, or Adolescence,” it is now classified as an anxiety disorder. The core features remain mostly unchanged, although the wording of the criteria has been modified to more adequately represent the expression of separation anxiety symptoms in adulthood. For example, attachment figures may include the children of adults with separation anxiety disorder, and avoidance behaviors may occur in the workplace as well as at school. Also, in contrast to DSM-IV, the diagnostic criteria no longer specify that age at onset must be before 18 years, because a substantial number of adults report onset of separation anxiety after age 18. Also, a duration criterion—“typically lasting for 6 months or more”—has been added for adults to minimize overdiagnosis of transient fears.

### **Selective Mutism**

In DSM-IV, selective mutism was classified in the section “Disorders Usually First Diagnosed in Infancy, Childhood, or Adolescence.” It is now classified as an anxiety disorder, given that a large majority of children with selective mutism are anxious. The diagnostic criteria are largely unchanged from DSM-IV.

## **Obsessive-Compulsive and Related Disorders**

The chapter on obsessive-compulsive and related disorders, which is new in DSM-5, reflects the increasing evidence that these disorders are related to one another in terms of a range of diagnostic validators, as well as the clinical utility of grouping these disorders in the same chapter. New disorders include hoarding disorder, excoriation (skin-picking) disorder, substance-/medication-induced obsessive-compulsive and related disorder, and obsessive-compulsive and related disorder due to another medical condition. The DSM-IV diagnosis of trichotillomania is now termed trichotillomania (hair-pulling disorder) and has been moved from a DSM-IV classification of impulse-control disorders not elsewhere classified to obsessive-compulsive and related disorders in DSM-5.

### **Specifiers for Obsessive-Compulsive and Related Disorders**

The “with poor insight” specifier for obsessive-compulsive disorder has been refined in DSM-5 to allow a distinction between individuals with good or fair insight, poor insight, and “absent insight/delusional” obsessive-compulsive disorder beliefs (i.e., complete conviction that obsessive-compulsive disorder beliefs are true). Analogous “insight” specifiers have been included for body dysmorphic disorder and hoarding disorder. These specifiers are intended to improve differential diagnosis by emphasizing that individuals with these two disorders may present with a range of insight into their disorder-related beliefs, including absent insight/delusional symptoms. This change also emphasizes that the presence of absent insight/delusional beliefs warrants a diagnosis of the relevant obsessive-compulsive or related disorder, rather than a schizophrenia spectrum and other psychotic disorder. The “tic-related” specifier for obsessive-compulsive disorder reflects a growing literature on the diagnostic validity and clinical utility of identifying individuals with a current or past comorbid tic disorder, because this comorbidity may have important clinical implications.

### **Body Dysmorphic Disorder**

For DSM-5 body dysmorphic disorder, a diagnostic criterion describing repetitive behaviors or mental

acts in response to preoccupations with perceived defects or flaws in physical appearance has been added, consistent with data indicating the prevalence and importance of this symptom. A “with muscle dysmorphia” specifier has been added to reflect a growing literature on the diagnostic validity and clinical utility of making this distinction in individuals with body dysmorphic disorder. The delusional variant of body dysmorphic disorder (which identifies individuals who are completely convinced that their perceived defects or flaws are truly abnormal appearing) is no longer coded as both delusional disorder, somatic type, and body dysmorphic disorder; in DSM-5 this presentation is designated only as body dysmorphic disorder with the absent insight/delusional beliefs specifier.

### **Hoarding Disorder**

Hoarding disorder is a new diagnosis in DSM-5. DSM-IV lists hoarding as one of the possible symptoms of obsessive-compulsive personality disorder and notes that extreme hoarding may occur in obsessive-compulsive disorder. However, available data do not indicate that hoarding is a variant of obsessive-compulsive disorder or another mental disorder. Instead, there is evidence for the diagnostic validity and clinical utility of a separate diagnosis of hoarding disorder, which reflects persistent difficulty discarding or parting with possessions due to a perceived need to save the items and distress associated with discarding them. Hoarding disorder may have unique neurobiological correlates, is associated with significant impairment, and may respond to clinical intervention.

### **Trichotillomania (Hair-Pulling Disorder)**

Trichotillomania was included in DSM-IV, although “hair-pulling disorder” has been added parenthetically to the disorder’s name in DSM-5.

### **Excoriation (Skin-Picking) Disorder**

Excoriation (skin-picking) disorder is newly added to DSM-5, with strong evidence for its diagnostic validity and clinical utility.

### **Substance/Medication-Induced Obsessive-Compulsive and Related Disorder and Obsessive-Compulsive and Related Disorder Due to Another Medical Condition**

DSM-IV included a specifier “with obsessive-compulsive symptoms” in the diagnoses of anxiety disorders due to a general medical condition and substance-induced anxiety disorders. Given that obsessive-compulsive and related disorders are now a distinct category, DSM-5 includes new categories for substance-/medication-induced obsessive-compulsive and related disorder and for obsessive-compulsive and related disorder due to another medical condition. This change is consistent with the intent of DSM-IV, and it reflects the recognition that substances, medications, and medical conditions can present with symptoms similar to primary obsessive-compulsive and related disorders.

### **Other Specified and Unspecified Obsessive-Compulsive and Related Disorders**

DSM-5 includes the diagnoses other specified obsessive-compulsive and related disorder, which can include conditions such as body-focused repetitive behavior disorder and obsessional jealousy, or unspecified obsessive-compulsive and related disorder. Body-focused repetitive behavior disorder is characterized by recurrent behaviors other than hair pulling and skin picking (e.g., nail biting, lip biting, cheek chewing) and repeated attempts to decrease or stop the behaviors. Obsessional jealousy is characterized by nondelusional preoccupation with a partner’s perceived infidelity.

## Trauma- and Stressor-Related Disorders

### Acute Stress Disorder

In DSM-5, the stressor criterion (Criterion A) for acute stress disorder is changed from DSM-IV. The criterion requires being explicit as to whether qualifying traumatic events were experienced directly, witnessed, or experienced indirectly. Also, the DSM-IV Criterion A2 regarding the subjective reaction to the traumatic event (e.g., “the person’s response involved intense fear, helplessness, or horror”) has been eliminated. Based on evidence that acute posttraumatic reactions are very heterogeneous and that DSM-IV’s emphasis on dissociative symptoms is overly restrictive, individuals may meet diagnostic criteria in DSM-5 for acute stress disorder if they exhibit any 9 of 14 listed symptoms in these categories: intrusion, negative mood, dissociation, avoidance, and arousal.

### Adjustment Disorders

In DSM-5, adjustment disorders are reconceptualized as a heterogeneous array of stress-response syndromes that occur after exposure to a distressing (traumatic or nontraumatic) event, rather than as a residual category for individuals who exhibit clinically significant distress without meeting criteria for a more discrete disorder (as in DSM-IV). DSM-IV subtypes marked by depressed mood, anxious symptoms, or disturbances in conduct have been retained, unchanged.

### Posttraumatic Stress Disorder

DSM-5 criteria for posttraumatic stress disorder differ significantly from those in DSM-IV. As described previously for acute stress disorder, the stressor criterion (Criterion A) is more explicit with regard to how an individual experienced “traumatic” events. Also, Criterion A2 (subjective reaction) has been eliminated. Whereas there were three major symptom clusters in DSM-IV—reexperiencing, avoidance/numbing, and arousal—there are now four symptom clusters in DSM-5, because the avoidance/numbing cluster is divided into two distinct clusters: avoidance and persistent negative alterations in cognitions and mood. This latter category, which retains most of the DSM-IV numbing symptoms, also includes new or reconceptualized symptoms, such as persistent negative emotional states. The final cluster—alterations in arousal and reactivity—retains most of the DSM-IV arousal symptoms. It also includes irritable or aggressive behavior and reckless or self-destructive behavior. Posttraumatic stress disorder is now developmentally sensitive in that diagnostic thresholds have been lowered for children and adolescents. Furthermore, separate criteria have been added for children age 6 years or younger with this disorder.

### Reactive Attachment Disorder

The DSM-IV childhood diagnosis reactive attachment disorder had two subtypes: emotionally withdrawn/inhibited and indiscriminately social/disinhibited. In DSM-5, these subtypes are defined as distinct disorders: reactive attachment disorder and disinhibited social engagement disorder. Both of these disorders are the result of social neglect or other situations that limit a young child’s opportunity to form selective attachments. Although sharing this etiological pathway, the two disorders differ in important ways. Because of dampened positive affect, reactive attachment disorder more closely resembles internalizing disorders; it is essentially equivalent to a lack of or incompletely formed preferred attachments to caregiving adults. In contrast, disinhibited social engagement disorder more closely resembles ADHD; it may occur in children who do not necessarily lack attachments and may have established or even secure attachments. The two disorders differ in other important ways, including correlates, course, and response to intervention, and for these reasons are considered separate disorders.

## Dissociative Disorders

Major changes in dissociative disorders in DSM-5 include the following: 1) derealization is included in the name and symptom structure of what previously was called depersonalization disorder and is now called *depersonalization/derealization disorder*, 2) dissociative fugue is now a specifier of dissociative amnesia rather than a separate diagnosis, and 3) the criteria for dissociative identity disorder have been changed to indicate that symptoms of disruption of identity may be reported as well as observed, and that gaps in the recall of events may occur for everyday and not just traumatic events. Also, experiences of pathological possession in some cultures are included in the description of identity disruption.

### Dissociative Identity Disorder

Several changes to the criteria for dissociative identity disorder have been made in DSM-5. First, Criterion A has been expanded to include certain possession-form phenomena and functional neurological symptoms to account for more diverse presentations of the disorder. Second, Criterion A now specifically states that transitions in identity may be observable by others or self-reported. Third, according to Criterion B, individuals with dissociative identity disorder may have recurrent gaps in recall for everyday events, not just for traumatic experiences. Other text modifications clarify the nature and course of identity disruptions.

## Somatic Symptom and Related Disorders

In DSM-5, somatoform disorders are now referred to as somatic symptom and related disorders. In DSM-IV, there was significant overlap across the somatoform disorders and a lack of clarity about their boundaries. These disorders are primarily seen in medical settings, and nonpsychiatric physicians found the DSM-IV somatoform diagnoses problematic to use. The DSM-5 classification reduces the number of these disorders and subcategories to avoid problematic overlap. Diagnoses of somatization disorder, hypochondriasis, pain disorder, and undifferentiated somatoform disorder have been removed.

### Somatic Symptom Disorder

DSM-5 better recognizes the complexity of the interface between psychiatry and medicine. Individuals with somatic symptoms plus abnormal thoughts, feelings, and behaviors *may or may not* have a diagnosed medical condition. The relationship between somatic symptoms and psychopathology exists along a spectrum, and the arbitrarily high symptom count required for DSM-IV somatization disorder did not accommodate this spectrum. The diagnosis of somatization disorder was essentially based on a long and complex symptom count of medically unexplained symptoms. Individuals previously diagnosed with somatization disorder will usually meet DSM-5 criteria for somatic symptom disorder, but only if they have the maladaptive thoughts, feelings, and behaviors that define the disorder, in addition to their somatic symptoms.

In DSM-IV, the diagnosis undifferentiated somatoform disorder had been created in recognition that somatization disorder would only describe a small minority of “somatizing” individuals, but this disorder did not prove to be a useful clinical diagnosis. Because the distinction between somatization disorder and undifferentiated somatoform disorder was arbitrary, they are merged in DSM-5 under somatic symptom disorder, and no specific number of somatic symptoms is required.

### Medically Unexplained Symptoms

DSM-IV criteria overemphasized the importance of an absence of a medical explanation for the somatic symptoms. Unexplained symptoms are present to various degrees, particularly in conversion disorder,

but somatic symptom disorders can also accompany diagnosed medical disorders. The reliability of medically unexplained symptoms is limited, and grounding a diagnosis on the absence of an explanation is problematic and reinforces mind-body dualism. The DSM-5 classification defines disorders on the basis of positive symptoms (i.e., distressing somatic symptoms plus abnormal thoughts, feelings, and behaviors in response to these symptoms). Medically unexplained symptoms do remain a key feature in conversion disorder and pseudocyesis because it is possible to demonstrate definitively in such disorders that the symptoms are not consistent with medical pathophysiology.

### **Hypochondriasis and Illness Anxiety Disorder**

Hypochondriasis has been eliminated as a disorder, in part because the name was perceived as pejorative and not conducive to an effective therapeutic relationship. Most individuals who would previously have been diagnosed with hypochondriasis have significant somatic symptoms in addition to their high health anxiety, and would now receive a DSM-5 diagnosis of somatic symptom disorder. In DSM-5, individuals with high health anxiety without somatic symptoms would receive a diagnosis of illness anxiety disorder (unless their health anxiety was better explained by a primary anxiety disorder, such as generalized anxiety disorder).

### **Pain Disorder**

DSM-5 takes a different approach to the important clinical realm of individuals with pain. In DSM-IV, the pain disorder diagnoses assume that some pains are associated solely with psychological factors, some with medical diseases or injuries, and some with both. There is a lack of evidence that such distinctions can be made with reliability and validity, and a large body of research has demonstrated that psychological factors influence all forms of pain. Most individuals with chronic pain attribute their pain to a combination of factors, including somatic, psychological, and environmental influences. In DSM-5, some individuals with chronic pain would be appropriately diagnosed as having somatic symptom disorder, with predominant pain. For others, psychological factors affecting other medical conditions or an adjustment disorder would be more appropriate.

### **Psychological Factors Affecting Other Medical Conditions and Factitious Disorder**

Psychological factors affecting other medical conditions is a new mental disorder in DSM-5, having formerly been included in the DSM-IV chapter “Other Conditions That May Be a Focus of Clinical Attention.” This disorder and factitious disorder are placed among the somatic symptom and related disorders because somatic symptoms are predominant in both disorders, and both are most often encountered in medical settings. The variants of psychological factors affecting other medical conditions are removed in favor of the stem diagnosis.

### **Conversion Disorder (Functional Neurological Symptom Disorder)**

Criteria for conversion disorder (functional neurological symptom disorder) are modified to emphasize the essential importance of the neurological examination, and in recognition that relevant psychological factors may not be demonstrable at the time of diagnosis.

## **Feeding and Eating Disorders**

In DSM-5, the feeding and eating disorders include several disorders included in DSM-IV as feeding and eating disorders of infancy or early childhood in the chapter “Disorders Usually First Diagnosed in Infancy, Childhood, or Adolescence.” In addition, brief descriptions and preliminary diagnostic criteria are

provided for several conditions under other specified feeding and eating disorder; insufficient information about these conditions is currently available to document their clinical characteristics and validity or to provide definitive diagnostic criteria.

### **Pica and Rumination Disorder**

The DSM-IV criteria for pica and for rumination disorder have been revised for clarity and to indicate that the diagnoses can be made for individuals of any age.

### **Avoidant/Restrictive Food Intake Disorder**

DSM-IV feeding disorder of infancy or early childhood has been renamed avoidant/restrictive food intake disorder, and the criteria have been significantly expanded. The DSM-IV disorder was rarely used, and limited information is available on the characteristics, course, and outcome of children with this disorder. Additionally, a large number of individuals, primarily but not exclusively children and adolescents, substantially restrict their food intake and experience significant associated physiological or psychosocial problems but do not meet criteria for any DSM-IV eating disorder. Avoidant/restrictive food intake disorder is a broad category intended to capture this range of presentations.

### **Anorexia Nervosa**

The core diagnostic criteria for anorexia nervosa are conceptually unchanged from DSM-IV with one exception: the requirement for amenorrhea has been eliminated. In DSM-IV, this requirement was waived in a number of situations (e.g., for males, for females taking contraceptives). In addition, the clinical characteristics and course of females meeting all DSM-IV criteria for anorexia nervosa except amenorrhea closely resemble those of females meeting all DSM-IV criteria. As in DSM-IV, individuals with this disorder are required by Criterion A to be at a significantly low body weight for their developmental stage. The wording of the criterion has been changed for clarity, and guidance regarding how to judge whether an individual is at or below a significantly low weight is now provided in the text. In DSM-5, Criterion B is expanded to include not only overtly expressed fear of weight gain but also persistent behavior that interferes with weight gain.

### **Bulimia Nervosa**

The only change to the DSM-IV criteria for bulimia nervosa is a reduction in the required minimum average frequency of binge eating and inappropriate compensatory behavior frequency from twice to once weekly. The clinical characteristics and outcome of individuals meeting this slightly lower threshold are similar to those meeting the DSM-IV criterion.

### **Binge-Eating Disorder**

Extensive research followed the promulgation of preliminary criteria for binge eating disorder in Appendix B of DSM-IV, and findings supported the clinical utility and validity of binge-eating disorder. The only significant difference from the preliminary DSM-IV criteria is that the minimum average frequency of binge eating required for diagnosis has been changed from at least twice weekly for 6 months to at least once weekly over the last 3 months, which is identical to the DSM-5 frequency criterion for bulimia nervosa.

### **Elimination Disorders**

No significant changes have been made to the elimination disorders diagnostic class from DSM-IV to DSM-5. The disorders in this chapter were previously classified under disorders usually first diagnosed in infancy, childhood, or adolescence in DSM-IV and exist now as an independent classification in DSM-5.

## Sleep-Wake Disorders

Because of the DSM-5 mandate for concurrent specification of coexisting conditions (medical and mental), sleep disorders related to another mental disorder and sleep disorder related to a general medical condition have been removed from DSM-5, and greater specification of coexisting conditions is provided for each sleep-wake disorder. This change underscores that the individual has a sleep disorder warranting independent clinical attention, in addition to any medical and mental disorders that are also present, and acknowledges the bidirectional and interactive effects between sleep disorders and coexisting medical and mental disorders. This reconceptualization reflects a paradigm shift that is widely accepted in the field of sleep disorders medicine. It moves away from making causal attributions between coexisting disorders. Any additional relevant information from the prior diagnostic categories of sleep disorder related to another mental disorder and sleep disorder related to another medical condition has been integrated into the other sleep-wake disorders where appropriate.

Consequently, in DSM-5, the diagnosis of primary insomnia has been renamed insomnia disorder to avoid the differentiation of primary and secondary insomnia. DSM-5 also distinguishes narcolepsy, which is now known to be associated with hypocretin deficiency, from other forms of hypersomnolence. These changes are warranted by neurobiological and genetic evidence validating this reorganization. Finally, throughout the DSM-5 classification of sleep-wake disorders, pediatric and developmental criteria and text are integrated where existing science and considerations of clinical utility support such integration. This developmental perspective encompasses age-dependent variations in clinical presentation.

### Breathing-Related Sleep Disorders

In DSM-5, breathing-related sleep disorders are divided into three relatively distinct disorders: obstructive sleep apnea hypopnea, central sleep apnea, and sleep-related hypoventilation. This change reflects the growing understanding of pathophysiology in the genesis of these disorders and, furthermore, has relevance to treatment planning.

### Circadian Rhythm Sleep-Wake Disorders

The subtypes of circadian rhythm sleep-wake disorders have been expanded to include advanced sleep phase syndrome, irregular sleep-wake type, and non-24-hour sleep-wake type, whereas the jet lag type has been removed.

### Rapid Eye Movement Sleep Behavior Disorder and Restless Legs Syndrome

The use of DSM-IV “not otherwise specified” diagnoses has been reduced by designating rapid eye movement sleep behavior disorder and restless legs syndrome as independent disorders. In DSM-IV, both were included under dyssomnia not otherwise specified. Their full diagnostic status is supported by research evidence.

DSM is the manual used by clinicians and researchers to diagnose and classify mental disorders. The American Psychiatric Association (APA) will publish DSM-5 in 2013, culminating a 14-year revision process. For more information, go to [www.DSM5.org](http://www.DSM5.org).

APA is a national medical specialty society whose more than 36,000 physician members specialize in the diagnosis, treatment, prevention and research of mental illnesses, including substance use disorders. Visit the APA at [www.psychiatry.org](http://www.psychiatry.org) and [www.healthyminds.org](http://www.healthyminds.org). For more information, please contact Eve Herold at 703-907-8640 or [press@psych.org](mailto:press@psych.org).